

The 4-Day Wave as Observed from the Upper Atmosphere Research Satellite Microwave Limb Sounder

1). R. Allen and J. L. Stanford

Department of Physics and Astronomy, Iowa State University, Ames, Iowa

L. S. Elson, E. F. Fishbein, L. Froidevaux and J. W. Waters

Jet Propulsion Laboratory/California Institute of Technology, Pasadena, California

Received _____; accepted _____

Short title: THE 4-DAY WAVE IN MLS DATA

Abstract.

The “4-day wave” is an eastward moving quasi-nondispersive feature with period near 4 days occurring near the winter polar stratopause. This paper presents evidence of the 4-day feature in Microwave Limb Sounder (MLS) temperature, geopotential height and ozone data from the late Southern winters of 1992 and 1993. Space-time spectral analyses reveal a double-peaked temperature structure consisting of one peak near the stratopause and another in the lower mesosphere, with an out-of-phase relationship between the two peaks. This double-peaked structure is reminiscent of recent three-dimensional barotropic/baroclinic instability model predictions, and is observed here for the first time. The height variation of the 4-day ozone signal is shown to compare well with a linear advective-photochemical tracer model. Negative regions of quasi-geostrophic potential vorticity (PV) gradient and positive Eliassen-Palm flux divergence are shown to occur, consistent with instability dynamics playing a role in wave forcing. Spectral analyses of PV derived from MLS geopotential height fields reveal a 4-day signal peaking near the polar stratopause. The three-dimensional structure of the 4-day wave resembles the potential vorticity “charge” concept, wherein a PV anomaly in the atmosphere (analogous to an electrical charge in a dielectric material) induces a geopotential field, a vertically oriented temperature dipole and circulation about the vertical axis.

1.. Introduction

The 4-day wave is a ubiquitous feature in the polar winter upper stratosphere. It was first observed in temperature data as a strong 4-day signal in zonal wavenumber 1 (Venne and Stanford 1979). Later studies showed that it consists of waves 1 through at least 4 all moving with the same phase speed, such that the period of wave 1 is near 4 days. Synoptic plots of temperature reveal a quasi-nondispersive "warm pool" of air which rotates eastward around the winter pole near the stratopause with a 4-day rotation period (Prata 1984; Lait and Stanford 1988b; Lawrence et al. 1995). Numerous observational (Venne and Stanford 1979, 1982; Prata 1984; Lait and Stanford 1988b; Randel and Lait 1991; Manney 1991; Fraser et al. 1993; Lawrence et al. 1995; Lawrence and Randel 1996) and theoretical (Hartmann 1983; Manney et al. 1988; Manney 1991; Manney and Randel 1993) studies have examined the 4-day phenomenon.

The wave has been observed in both hemispheres, but is stronger in the Southern Hemisphere (Venne and Stanford 1982). The horizontal phase structure is variable, with both poleward and equatorward momentum flux events (Venne and Stanford 1982; Prata 1984; Manney 1991). The wave is generally barotropic, although some episodes show strong equatorward heat flux (Randel and Lait 1991). The episodes studied by Randel and Lait (1991) and Lawrence and Randel (1996) reveal a dipole structure of positive and negative Eliassen-Palm flux divergence associated with the 4-day wave, with the positive region overlapping negative quasi-geostrophic potential vorticity gradient, a signature of instability processes.

Theoretical studies have focused on instability of the jet structure as the mechanism for 4-day wave growth. Hartmann (1983) and Manney et al. (1988) showed that periods of 4-day wave growth are consistent with barotropic instability of the stratospheric polar night jet. Manney and Randel (1993) used a 3-D instability model to show that realistic 4-day wave growth rates exist in monthly mean background wind states only if both barotropic and baroclinic components are included. Their model also predicted

a double-lobed structure in the perturbation temperature field with one lobe in the upper stratosphere and another in the lower mesosphere, with rapid phase variation in between.

The present paper takes advantage of the coverage and vertical resolution of the Microwave Limb Sounder (MLS) on board the Upper Atmosphere Research Satellite (UARS) to examine the 4-day wave during the late austral winters 1992 and 1993. As will be shown below, strong 4-day signals exist in MLS temperature, geopotential height and ozone. The vertical coverage provided by MLS allows resolution of a double-peaked structure in perturbation temperature with an out-of-phase relationship between the upper and lower peaks, as predicted by Manney and Randel (1993). Spectral analyses of MLS ozone reveal a 4-day signal in zonal wavenumber 1 near the polar stratopause. A linear advective-photochemical model is used to calculate the ozone response to temperature and wind perturbations associated with the 4-day wave. The vertical variation of model ozone amplitude is in good agreement with observations of MLS ozone. Negative regions of quasi-geostrophic potential vorticity ($\nabla \cdot \mathbf{V}$) gradient and positive Eliassen-Palm flux divergence are shown to occur, consistent with wave forcing by instability dynamics.

The 4-day feature is also shown to exist in quasi-geostrophic PV derived from MLS geopotential heights. The three-dimensional structure of this signal resembles the PV "charge" discussed by Hoskins et al. (1985) and more recently by Bishop and Thorpe (1994). The latter authors have also extended the concept to Ertel PV (Thorpe and Bishop 1995), although the present paper focuses on quasi-geostrophic PV. In this electrostatics analogy, a PV anomaly in the atmosphere acts like an electrical charge in a dielectric material, with an associated vector field (independent of static stability, density and boundary conditions) that produces "action-at-a-distance". Isolated PV charges induce geopotential anomalies, circulation about the vertical axis and a vertically oriented temperature dipole, similar to observations of the 4-day wave presented in this

study.

2. Microwave Limb Sounder Data

a. The MLS instrument

The Microwave Limb Sounder is one of ten instruments on board the Upper Atmosphere Research Satellite. MLS uses a 1.6 m scanning antenna to observe the atmospheric limb emission simultaneously in spectral bands at 63, 183 and 205 GHz (Barath et al. 1993). These observations allow the determination of chlorine monoxide, ozone, water vapor, nitric acid, sulfur dioxide, temperature and geopotential height in the stratosphere and mesosphere during both day and night and in the presence of stratospheric clouds and aerosols. This study analyzes temperature, geopotential height and ozone data from two austral late-winter time periods, 14 August – 20 September 1992 and 9 August - 16 September 1992, when MLS was viewing southward (coverage from 80 S to 34 N).

b. Temperature Data

MLS temperature is obtained by viewing limb emission of O_2 at 63 GHz. The version 3 data (used here) are retrieved on every other UARS pressure level n (the UARS Level 3 pressure grid is $1000 \times 10^{-n/6}$ hPa, where $n = 0, 1, 2, \dots$) and have uncertainties ranging from 1.5 K at 22 hPa to 6.4 K at 0.46 hPa. In this paper MLS temperatures are used at 5 pressure levels: 22, 10, 4.6, 2.2, 1, 0.46, 0.22, 0.1 hPa. Results from the two highest levels, 0.22 and 0.1 hPa, should be viewed with some caution due to large errors which move the retrievals toward climatology. Validation information on version 3 MLS temperature is available in Fishbein et al. (1996).

c. Ozone Data

MLS ozone retrievals are obtained from limb emission in two bands centered near 205 and 183 GHz. The version 3 ozone data are retrieved on every other UARS pressure level. The 205 GHz data have smaller uncertainties throughout the stratosphere, while the 183 GHz data are less noise-limited in the mesosphere. Both 205 and 183 GHz ozone data are used in this paper from 22 to 0.22 hPa. Validation information on version 3 MLS ozone is available in Froidevaux et al. (1996) and Ricaud et al. (1996). Single-profile precision for ozone retrievals is in the 0.1 to 0.4 ppmv range, depending on altitude.

d. Geopotential Height Data

The MLS geopotential height data set used in this study is a preliminary data set which is not available in version 3, but is planned for future versions. The geopotential height is obtained by measuring pressure as a function of tangent-point height. Pressure is retrieved from the same set of measurements as temperature, while tangent-point height is determined from an encoder mounted on the antenna axis and UARS orbit/attitude knowledge. The uncertainty in pressure can be represented in terms of the expected tangent point height. The uncertainty ranges from 110 m at 10 hPa to 40 m at 0.46 hPa, thus also giving an approximation to the errors in the geopotential height data. Uncertainties in a coordinate transformation involved in the geopotential height retrieval is known to cause significant errors in the calculation of the zonal mean and high-frequency wave components (periods less than about 2.5 days). However, the analyses performed in this paper (zonal wave 1, 3-5 day periods) have sufficiently long periods to avoid significant effects from these errors. Good agreement is found on these spatial and temporal scales between MLS temperature (coined $T_{retrieved}$ in this paper) and temperature derived from MLS geopotential height data (T_{geoh}), as discussed in Section 4a.

3. Analysis Procedure

MIS data used in this analysis were produced by the asymptotic mapping method (Salby 1982a,b; Lait and Stanford 1988a). The specific technique used for this study is presented in detail in Elson and Froidevaux (1993). Briefly, data are binned by latitude and separated into ascending and descending time series. The time and space axes are rotated to a new coordinate system where a Fast Fourier Transform (FFT) is applied to calculate the spectral coefficients. An inverse transformation and interpolation allows data to be mapped on synoptic maps which are processed every 12 hours. Satellite orbital period drift generally limits the procedure to approximately one week analysis periods. The procedure here is essentially a re-mapping in time so that periods longer than a week can be analyzed. Data are interpolated onto horizontal grids of 4 degrees latitude by 5 degrees longitude and onto the standard UARS pressure levels.

The twice-daily synoptic maps described above were decomposed in space and time by applying a two-dimensional sinusoidal transform (see appendix of Ziemke and Stanford 1990). Power spectral density is defined in this study as $(2\Delta f)^{-1}[A^2(k, \omega) + B^2(k, \omega)]$, and amplitude is defined as $[A^2(k, \omega) + B^2(k, \omega)]^{0.5}$, where Δf is the unit bandwidth $(NDAT \cdot \Delta t)^{-1}$. $NDAT$ is the number of data points in the time series ($NDAT=38$ for 1992, $NDAT=30$ for 1993) and Δt is the temporal sampling interval (0.5 day). $A(k, \omega)$ and $B(k, \omega)$ are the cosine and sine coefficients of the discrete Fourier expansions

$$\{T_{\pm}, \Phi_{\pm}, \mu_{\pm}\}(x, t) = \sum_{k=0}^{\pi/\Delta x} \sum_{\omega=0}^{\pi/\Delta t} [A_{\pm}(k, \omega) \cdot \cos(kx \pm \omega t) + B_{\pm}(k, \omega) \cdot \sin(kx \pm \omega t)]. \quad (1)$$

Here T , Φ and μ are temperature, geopotential and ozone mixing ratio; \pm ($-$) refers to westward (eastward) components; t is time; x is zonal distance; Δx is the zonal grid interval of $\frac{2\pi a \cdot \cos \phi}{72}$; k is the zonal wavenumber; ω is the angular frequency; a is the earth's radius, and ϕ is latitude. A 1-2-1 running mean was applied in frequency space to power spectral density, but not to amplitude. The time mean was removed before applying the FFT to focus the analysis on traveling waves.

Filtered time series in this study were produced by first applying the band-pass filter response function (with half amplitude points at 3 and 5 day periods) given by Murakami (1979) to the temporal FFT coefficients and then recombining the coefficients through an inverse FFT.

As a comparison to the MLS temperature measurements, temperature was also calculated from MLS geopotential height using the hydrostatic approximation $T = R^{-1} \partial \Phi / \partial (\ln p)$ with $R = 287 \text{ J kg}^{-1} \text{ K}^{-1}$ (dry air gas constant) and constant vertical spacing $d(\ln p) = 0.357$ (determined from UARS standard pressure surfaces). Quasi-geostrophic potential vorticity, q' , and potential vorticity gradient, \bar{q}_y , were calculated using the spherical formulation of Matsuno (1970):

$$q' = \frac{1}{f a^2} \left[\frac{\Phi'_{\lambda\lambda}}{\cos^2 \phi} + \frac{f^2}{\cos \phi} \left(\frac{\cos \phi}{f^2} \Phi'_{\phi} \right)_{\phi} + \frac{f^2 a^2}{\rho_o} \left(\frac{\rho_o \Phi'_z}{N^2} \right)_z \right] \quad (2)$$

$$\bar{q}_y = \frac{2\Omega \cos \phi}{a} - \left[\frac{(\bar{u} \cos \phi)_{\phi}}{a^2 \cos \phi} \right]_{\phi} - \frac{1}{\rho_o} \left(\frac{\rho_o f^2}{N^2} \bar{u}_z \right)_z \quad (3)$$

Here u is the zonal wind; $\rho_o = \rho_{oo} e^{-z/H}$; ρ_{oo} is the density at the surface (1000 hPa); z is the log-pressure coordinate; λ is longitude; f is the Coriolis parameter, and Ω is the earth's angular frequency. Overbars denote zonal means and primes denote deviation from the zonal mean. Subscripts denote partial derivatives. The Brunt-Väisälä parameter and atmospheric scale height were given the constant values $N^2 = 4 \times 10^{-4} \text{ s}^{-2}$ and $H = 7 \text{ km}$. q' was calculated from MLS geopotential height data, while \bar{q}_y was calculated from United Kingdom Meteorological Office (UKMO) zonal winds. [MLS data were not used in the calculation of \bar{q}_y due to the known inaccuracy in the zonal mean geopotential height (see Section 2d).

UKMO horizontal winds and temperatures on UARS pressure surfaces, obtained from the Goddard Space Flight Center (GSFC) Distributed Active Archive Center (DAAC), were interpolated with a cubic spline onto horizontal grids of 4 degrees latitude by 5 degrees longitude. These winds and temperatures were used in the calculation of \bar{q}_y and also in the tracer model discussed in Section 4c.

The quasi-geostrophic Eliassen-Palm (EP) flux vector and the EP flux divergence were calculated with the the following spherical formulation:

$$\mathbf{F} = (F_\phi, F_z) = \rho_o a \cos \phi \left(-\overline{u'v'}, \frac{R}{H} f \frac{\overline{v'T'}}{N^2} \right) \quad (4)$$

$$\nabla \cdot \mathbf{F} = \frac{1}{a \cos \phi} \frac{\partial}{\partial \phi} (\cos \phi F_\phi) + \frac{\partial F_z}{\partial z}. \quad (5)$$

Here u' and v' (meridional wind) are calculated from MLS geopotential heights (using the geostrophic approximation) and T' is from MLS retrieved temperatures. In the EP flux diagram (Fig. 11) the vector components are multiplied by $2\pi a \cos \phi$. This factor accounts for the spherical geometry of the earth to make the arrow pattern look non-divergent if and only if $\nabla \cdot \mathbf{F} = 0$ (see Edmon et al. 1980).

4. Results

a. Temperature

Space-time spectral decomposition is a useful tool for isolating the 4-day wave from other atmospheric signals. Figure 1a presents zonal wavenumber 1 power spectral density (defined above Eq. 1 and hereafter *power*) for MLS retrieved temperature (hereafter $T_{retrieved}$) at 72° South latitude for 18 August - 6 September 1992. A clear 4-day eastward period signal exists near 1 hPa (48 km) and a somewhat weaker signal occurs near 0.22 hPa (60 km). These signals are also evident in temperature derived from MLS geopotential height (hereafter T_{geoh}), Fig. 1 b, although the lower signal maximizes near 2.2 hPa (42 km). An even stronger 4-day signal is found during 2 - 16 September 1993 in both $T_{retrieved}$ (Fig. 1 c) and T_{geoh} (Fig. 1 d). The 4-day signal dominates the traveling wave spectra for this time period with two lobes centered near 2.2 hPa (42 km) and 0.22 hPa (60 km). Sections 4d and 4f discuss further the validity of the 4-day signal in terms of uncertainty and statistical significance.

The 1992 4-day wave event was further isolated by filtering a 38 day series of $T_{retrieved}$ for zonal wavenumber 1 and eastward periods of 3-5 days and plotting on

a Hovmöller (time vs. longitude) diagram for 64 S and 1 hPa (Fig. 2). The 4-day wave amplitude increases over the first 10 days and remains constant for another 10 days before dissipating near 1 September 1992. The mode makes at least 5 revolutions around the pole with a rotation period near 3.8 days,

Spectral plots of $T_{retrieved}$ for zonal wavenumbers 2 and 3 (not shown here) for the 1992 event show signals at 1 hPa with eastward periods of 2.0 and 1.3 days, respectively. These sinusoidal components move at the same phase speed of the wave 1, 4-day signal and combine with wave 1 to form a ‘(warm pool’ of air at 1 hPa which rotates around the pole with a period near 4 days. This feature is shown in Fig. 3, where $T_{retrieved}$ data have been filtered over 14 August - 20 September 1992 for waves 1 through 3 and eastward rotation periods of 3-5 days. (Wave 4 and higher were not analyzed because the frequency of interest, 4 days divided by integer wavenumber, is near or outside the Nyquist limit of approximately 1 day⁻¹.) The warm pool rotates around the pole as an identifiable entity throughout the record shown displaying the quasi-nondispersive (soliton-like) nature of the feature.

The perturbation temperature ($T_{retrieved}$) amplitude for wave 1 and 3.8 day eastward period is plotted in Fig. 4a for the 1992 4-day wave event. A large lobe exists in the upper stratosphere peaking near 48 km and 72 S with maximum amplitude of 4.3 K. A smaller upper lobe is centered at 60 km and 68 S. This double-lobed structure is similar to that predicted by the instability study of Manney and Randel (1993). In their results, the upper lobe is larger and extends well into the mesosphere. However, $T_{retrieved}$ is largely climatology at 66 km, forcing zonally asymmetric wave amplitudes to zero, therefore the abrupt fall-off with height above 60 km in Fig. 4a may not be real. The 1993 series also displays a double-lobed structure (Fig. 4b), but each lobe contains two maxima. The lower lobe: has maxima near 42 km, 76 S (amplitude of 5.3 K) and 48 km, 60 S, while the upper lobe is an extended region with two weak maxima at 60 km, 50 S and 60 km, 64 S.

Manney and Randel (1993) also predict rapid vertical phase variation between the two lobes. Such behavior can be explicitly seen in height vs. longitude plots of $T_{retrieved}$ at 72 S, filtered for waves 1 through 3 and eastward rotation periods of 3-5 days. Figure 5a presents 8 days during the peak amplitude of the 1992 time series. The plots reveal a warm anomaly near 48 km bounded above by a cold anomaly near 60 km. This dipole pattern continues throughout the 4-day wave life-cycle, suggesting that the two peaks constitute one entity. The 1993 series also reveals this vertical temperature dipole (Fig. 5b). Here the lower peak has a strong wave 1 structure which begins near 48 km and descends to 42 km over this 8 day period. The upper feature, which maximizes near 60 km, is approximately 180 degrees out of phase with the lower feature. The quadrupole pattern produced by the two out-of-phase lobes remains stable for at least two weeks, through the end of the UARS month (16 September 1993), when MLS changed to northward viewing (coverage from 80 N to 34 S).

b. Geopotential

The vertical temperature dipole structure seen in Fig. 5 suggests that the perturbation geopotential amplitude should increase with height throughout the region of the lower temperature lobe and decrease with height in the region of the upper lobe. This is confirmed in Fig. 6a, which shows the geopotential amplitude for the 1992 series for eastward moving wave 1 with a period of 3.8 days. The signal maximizes near 57 km, at the boundary between the warm and cold anomalies (see Fig. 5a), and decreases above. The 1993 geopotential amplitude (Fig. 6b) peaks near 48 km, consistent with the perturbation temperature structure for that series (Fig. 5b). The geopotential signal decreases with altitude more rapidly in the 1993 time series, due to a larger amplitude upper temperature lobe in 1993 than in 1992. The meridional structure shown here is similar to previous observations of the 4-day wave in geopotential height (Manney 1991; Lawrence and Randel 1996).

c.. ozone

Other than the unpublished conference report by Randelet al. (1992), the present paper is the first to present observations of the 4-day wave in ozone. East-west power spectra for wave 1MLS ozone (both 205 GHz and 183 GHz data) at 72 S during 18 August - 6 September 1992 are presented in Fig. 7. A 4-day peak exists near the stratopause (1.0 hPa), coincident with the $T_{retrieved}$ maximum at the same altitude, Fig. 1a. A 2-day eastward period signal exists at 2.2 hPa; the source of this signal is unknown and is not thought to be related to the 4-day wave. The 2-day feature does not exist in spectra from 5- 20 September 1992.

To ascertain the relationship between the 4-day feature in ozone and the other atmospheric variables, a linear advective-photochemical model was used to calculate the ozone response to temperature and meridional wind perturbations associated with the 4-day wave. The procedure combines the linearized thermodynamic and ozone continuity equations to obtain an expression for the ozone mixing ratio (Hartmann and Garcia 1979; Rood and Douglass 1985; Randel 1993). Because the 4-day wave amplitudes in the various components (T' , u , v and μ) are observed to be much less than their respective zonal mean values ($T' \ll \bar{T}$, etc.), we use a linearized thermodynamic equation,

$$T'_t + \bar{u}T'_x + v\bar{T}'_y + w'S = 0 \quad (6)$$

$$S = \frac{1}{H} [\kappa\bar{T} + H\bar{T}_z]$$

where w' is the vertical wind component, $\kappa = R/c_p \approx 2/7$ and $H = 7$ km, and a linearized ozone continuity equation,

$$\mu'_t + \bar{u}\mu'_x + v'\bar{\mu}_y + w'\bar{\mu}_z = -\Gamma\mu' - \Theta T'. \quad (7)$$

The photochemical relaxation frequency, Γ , and the linearized ozone response to small temperature perturbations, Θ , are shown in Fig. 8a, adapted from Hartmann and Garcia

(1979) who studied ozone transport by planetary waves in the winter stratosphere using a linearized β -plane model centered at 60° . Radiative damping of the temperature perturbations is neglected, since the radiative time scale is of order 10 days in the region of interest (26 to 57 km), see Hartmann and Garcia (1979), while the advective time scale $[k(\bar{u} - c)]^{-1}$ is of order 1-2 days from 26 to 42 km, and the photochemical time scale Γ^{-1} is less than 1 day from 42 to 57 km. Combining Eqs. 4 and 5 gives the following differential equation for μ' :

$$\left(\frac{\partial}{\partial t} + \bar{u}\frac{\partial}{\partial x} + \Gamma\right)\mu' + \left(\bar{\mu}_y - \frac{\bar{\mu}_z}{S}\bar{T}_y\right)v' - \left[\frac{\bar{\mu}_z}{S}\left(\frac{\partial}{\partial t} + \bar{u}\frac{\partial}{\partial x}\right) - \Theta\right]T' = 0. \quad (8)$$

The perturbation quantities are assumed to have zonally propagating sinusoidal solutions: $\mu' = \mu_A \cdot \cos[k(x - ct)] + \mu_B \cdot \sin[k(x - ct)]$. Similar dependence is assumed for v' and T' . Substituting into Eq. 8, the resulting expression for the ozone mixing ratio amplitude ($\mu_{amp}^2 = \mu_A^2 + \mu_B^2$, similarly for v_{amp} and T_{amp}) is

$$\mu_{amp}^2 = \frac{1}{(a^2 + \Gamma^2)} \left[(b^2 + \Theta^2) T_{amp}^2 + \xi^2 v_{amp}^2 + 2b\xi(v_B T_A - v_A T_B) + 2\Theta\xi(v_A T_A + v_B T_B) \right] \quad (9)$$

$$\xi \equiv \bar{\mu}_y - \bar{\mu}_z \bar{T}_y / S \quad a \equiv k(\bar{u} - c) \quad b \equiv k(\bar{u} - c) \bar{\mu}_z / S.$$

Figure 8b presents the resulting model ozone amplitude for the wave 1, 4.0 day period signal at 60 S (thick solid line). Also plotted are the observed ozone amplitudes at 60 S for 205 GHz (thin solid) and 183 GHz (dotted) data. Error bars on the observed data were calculated using the procedure described in Section 4d. Reasonable agreement is found between observed and model amplitudes, with low values in the middle stratosphere and lower mesosphere, and a maximum in the upper stratosphere (near 42 km). Results from a purely advective model ($\Gamma = \Theta = 0$) and a purely photochemical model ($\mu_{amp} = (\Theta/\Gamma) T_{amp}$, see Randel 1990) show a transition region between dynamical and photochemical control from approximately 37-49 km, where both dynamics and photochemistry play an important role.

The good agreement (at least to first order) between model and observations suggests that the observed 4-day signal in ozone is produced largely by the coupled effects of constituent advection by the 4-day wave perturbation winds in the presence of zonal mean constituent gradients and photochemical processes associated with the 4-day wave perturbation temperature and mixing ratio anomalies.

d. Uncertainty in 4-Day Wave Amplitude

In order to assess the uncertainty in the 4-day wave amplitude calculation, a random number generator was used to make a noise data set with Level3AT format (along-track, evenly spaced in time) with values between -0.5 and +0.5 (arbitrary units), giving a Root Mean Square (RMS) variation of 0.29. This represents the random noise in a given MLS measured quantity with an RMS error (defined here as uncertainty) of 0.29. These data were passed through the synoptic mapping routine described in Section 2 and synoptic maps were produced. The RMS variation in the synoptic maps is 0.19, which is 65% of the input value. The decrease is mainly due to interpolation to a regular horizontal grid. Fourier decomposition was then applied to isolate the amplitude of the wave 1, 3.5 day component from 500 different time series, each randomly generated. This Monte Carlo procedure yielded RMS variations in amplitude of 0.018 for the 1992 simulated series (38 time components used, 72 longitudes) and 0.020 for the 1993 simulated series (30 time components used, 72 longitudes). The uncertainty in amplitude is $0.020/0.29 \approx 1/16$ times the uncertainty in the Level3AT data for 1992 and $1/14$ for 1993. The difference between $1/16$ and $1/14$ is due to the different bandwidth used in each year. This method can now be used to obtain estimated uncertainties in 4-day wave amplitudes observed in MLS temperature and ozone.

The estimated uncertainty in Version 3 MLS temperature at 1 and 2.2 hPa is 3.4 and 2.2 K, respectively (Fishbein et al. 1996). The estimated 4-day wave amplitude uncertainty in the lower peak of Fig. 4a (1992) is $3.4/16 \approx 0.21$ K, which is much

smaller than the observed amplitude of 4.3 K. In 1993 the uncertainty in the lower peak (Fig. 4b) is $2.2/1.4 = 0.16$ K, which is much less than the observed amplitude of 5.2 K. This Monte Carlo analysis shows that the observed 4-day wave amplitudes are not significantly affected by random noise in the data.

This method was also used to obtain estimated errors for 205 and 183 GHz MLS ozone data. The error bars shown in Fig. 8b were computed using the above formulation along with data uncertainties at different pressure levels as given by Froidevaux et al, (1996).

c. Statistical Assessments

Because the 4-day wave is a well established atmospheric phenomenon (see references in introduction), statistical assessment of the 4-day features found here can be made on the basis of *a priori* statistics (Madden and Julian, 1971). Briefly, before analysis it is decided to examine spectral features near 4 day periods. (If spectral peaks are examined at previously unreported frequencies, *a posteriori* statistics must be used, Madden and Julian, 1971). The null hypothesis is made that any spectral peak at 4 days is due to random sampling fluctuations. Under this hypothesis, the ratio of spectral peak to nearby spectral background can be estimated statistically. For the Fourier transform spectra used here, chi squared statistics are employed to estimate the maximum peak to background ratio that will occur in 95% of the cases for random sampling fluctuations. If the observed peak exceeds this ratio, with 95% confidence the spectral feature can be said to be statistically significant.

Figures 9a,b show the $T_{retrieved}$ spectra at 2 and 0.2 hPa during 2- 16 September 1993. These are plotted without smoothing in Figs. 9a,b, and correspond to the smoothed plot shown in Fig. 1c. Also indicated by dashed lines in Fig. 9 are the estimated background and 95% confidence limit (for two degrees of freedom, appropriate to this case of unsmoothed Fourier transform spectral estimates). Both features easily

satisfy a *a priori* statistical significance.

The 1992 temperature spectral 4-day peak, shown smoothed in Fig. 1a, reveals a strong peak centered at 1 hPa which also easily exceeds the *a priori* 95% confidence limit (not shown here). The weaker temperature peak at 0.22 hPa is close to a stronger lower frequency feature. By examination of unsmoothed spectral results (not shown here) it can be shown that the 0.22 hPa 4-day peak also meets the criterion for *a priori* statistical significance.

The 183 GHz ozone spectrum at 1 hPa for 1992 (shown smoothed in Fig. 7b) is plotted unsmoothed in Fig. 9c. The 4-day peak exceeds the 95% confidence limit and is judged statistically significant. The 205 GHz ozone 4-day peak at 1 hPa (Fig. 7b) also exceeds the 95% confidence limit (not shown here).

f. Zonal Wind, \bar{q}_y and EP Flux Structure

Theoretical studies show that barotropic and/or baroclinic instability of the polar night jet can produce quasi-nondispersive modes similar to the observed 4-day wave. The instability episodes are generally accompanied by regions of negative quasi-geostrophic potential vorticity gradient, \bar{q}_y , and non-zero EP flux divergence. Figure 10a shows the zonal mean UKMO east-west wind averaged over the life-cycle of the 1992 4-day wave event (18 August - 6 September 1992). A strong stratospheric westerly jet is centered near 34 km, 60 S. The associated wind curvature produces a region of negative \bar{q}_y near 50 S at altitudes above 50 km (Fig. 10b). Daily wind maps reveal a “double-jet” structure near the beginning of the time series which is often present during 4-day wave events (Manney 1991; Randel and Lait 1991). Manney et al. (1996) showed that daily maps of Ertel PV (on isentropic surfaces) calculated from UKMO winds sometimes exhibit unrealistic structure in the Southern Hemisphere winter middle and upper stratosphere poleward of about 70 S. Our analyses here, however, focus on quasi-geostrophic PV averaged over 20 days. Comparison of Fig. 10a with National Meteorological Center

(NMC) zonal winds (not shown here) suggests good overall agreement. However, \bar{q}_y calculated from the NMC data (which extends only to 54 km) does not show the negative region near 50 S for this time-averaged period, although negative \bar{q}_y does exist near 50 S, 54 km for several of the individual NMC-derived daily maps during the first half of this time series.

Linear theory requires that wave growth by barotropic instability be accompanied by negative PV gradients. (The integral over latitude of \bar{q}_y times a positive definite quantity must be zero for wave growth to occur by *barotropic* instability. However, *baroclinic* instability is possible with positive \bar{q}_y , given suitable variation with latitude of the vertical gradient of \bar{q}_y at the lower boundary, see Holton 1992.) The presence here of negative \bar{q}_y regions is consistent with the fact that negative \bar{q}_y is a necessary, although not sufficient, condition for wave growth by barotropic instability.

Figure 11 displays the Eliassen-Palm (EP) flux diagram for this wave episode, giving details about the net effect on the zonal mean state of eddy and momentum fluxes due to wave events. The EP flux was calculated using MLS temperature and geopotential height from 18 August to 6 September 1992, filtered for wave 1 and 3.8 day eastward period. The geostrophic approximation was used to calculate the winds from the 4-day wave perturbation geopotential field. EP vectors are plotted over contours of $DP' = \mathbf{V} \cdot \mathbf{F} / (\rho_0 a \cos \phi)$ which is a force per unit mass on the zonal mean wind (in $\text{ms}^{-1} \text{day}^{-1}$) due to the eddy effects associated with the wave 1 component of the 4-day wave. Positive (negative) regions of DP' act to accelerate (decelerate) the zonal mean westerly wind. Positive DP' exists between 50 and 75 S at the top of the plot, near 57 km, acting to accelerate the lower mesospheric winds, while a negative DP' region near 54 km, 72 S acts to decelerate the stronger stratopause winds (see Fig. 10a). Figure 11 predicts eddy-induced reduction of vertical shear in the 60-75 S lower mesosphere region, consistent with baroclinic instability mechanisms driving the 4-day wave in the lower mesosphere. The EP flux structure above 48 km is similar to the model results

from Manney and Randel (1993, their Fig. 2d), with a region of negative DF bounded above and to the north by positive DF .

The EP flux vectors indicate strong equatorward heat flux (downward pointing arrows) and equatorward momentum flux (poleward pointing arrows) above 48 km and poleward of 60 S (also similar to Manney and Randel 1993, Fig. 2d) and largely poleward momentum flux elsewhere. The vectors are predominantly vertical (indicating baroclinic processes) in the lower mesosphere, while strong momentum flux (indicated by horizontal components of EP flux vectors) near 43 km, 60 S suggests barotropic processes play a sizable role there. This is confirmed by the positive (70 S) and negative (60 S) DF regions at 43 km, which act to reduce the latitudinal shear associated with the stratospheric jet (a signature of barotropic instability).

Manney and Randel (1993) and Lawrence and Randel (1996) show overlapping regions of negative \bar{q}_y and positive $\nabla \cdot \mathbf{F}$ during 4-day wave episodes. This spatial coincidence is recognized as a source region of wave activity $A = \frac{1}{2} \tilde{q}'^2 / \bar{q}_y$, where q' represents the perturbation quasi-geostrophic potential vorticity. Neglecting diabatic and frictional effects, a conservation equation can be written (Andrews and McIntyre (1976, 1978):

$$\frac{\partial A}{\partial t} + \nabla \cdot \mathbf{F} = 0. \quad (10)$$

Clearly, when \bar{q}_y is negative, the wave activity will increase in magnitude when $\nabla \cdot \mathbf{F} > 0$. One region of negative \bar{q}_y occurs in the vicinity of strong positive DF near 57 km, 52 S. This is consistent with instability dynamics driving the 4-day wave in the lower mesosphere. According to Eq. 10, wave activity will also increase if $\bar{q}_y > 0$ and $\nabla \cdot \mathbf{F} < 0$. The regions of negative DF near 54 km, 80 S to 65 S and 43 km, 60 S overlap positive \bar{q}_y , suggesting instability forcing may also exist in these regions.

g, Quasi-Geostrophic Potential Vorticity

Spectral analyses of quasi-geostrophic potential vorticity (PV) calculated from MLS geopotential heights reveal 4-day signals in both the 1992 and 1993 series. Figure 12 shows the east-west PV power spectra for 72 S during 18 August - 6 September 1992 (Fig. 12a) and 2-- 16 September 1993 (Fig. 12 b). Clear 4-day eastward period signals in PV are seen near the stratopause (48 km or 1 hPa) during both time periods. (In 1992, a significant longer-period feature is seen in the upper stratosphere. This eastward-moving power with periods of several weeks is not directly related to the 4-day wave considered here.) A height vs. latitude plot of PV amplitude for wave 1 and 3.8 day eastward period is provided in Fig. 13a for the 1992 series. A region of large PV amplitude is centered near 50 km, 68 S. Figure 13b shows the wave 1, 3.75 day signal in 1993, which has a similar structure to the 1992 feature, although centered slightly lower and with contours somewhat more compact.

The three-dimensional structure of the perturbation PV resembles the "PV charge" concept discussed by Hoskins et al. (1985) and more recently by Bishop and Thorpe (1994), who considered the effects on the atmosphere of a uniform spherical quasi-geostrophic PV anomaly in a basic state with constant static stability and density and in a two-layer vertical static stability basic state representing the effects of the tropopause. In this analogy, the atmosphere represents an isotropic dielectric material; static stability represents dielectric properties; geopotential (or streamfunction) represents electric potential, and an isolated PV anomaly represents an electrostatic charge, with an accompanying vector field that produces "action-at-a-distance" (a fundamental concept in PV thinking). This "PV field" (which is independent of static stability, density and boundary conditions) induces perturbation temperature, geopotential and wind fields as shown in Fig. 14a (for the case of constant static stability), adapted from Figs. 1 and 2 of Bishop and Thorpe (1994). Here a negative spherical PV charge induces a vertical temperature dipole with cold temperature perturbation on top, spherical geopotential

anomaly and anti-cyclonic circulation about the vertical axis.

In comparison, Fig. 14b shows the amplitude of the wave 1, 3.8 day period mode during the 1992 series as a function of height and latitude. The hatched region includes PV amplitude greater than 90% of the maximum. Solid lines are perturbation temperature amplitude of 60% to 90% of the maximum, incremented by 10%, and the dashed lines are perturbation geopotential amplitude at every 10% of the maximum. Here we see the PV anomaly bounded above and below by two temperature lobes, which are out-of-phase, resembling the cold and warm perturbations in Fig. 14a. The geopotential anomaly surrounds the region of large PV amplitude, although centered slightly higher. There would also be an induced circulation about the vertical axis associated with the geopotential anomaly, similar to that shown in Fig. 14a.

This analogy suggests that the 4-day wave may be more fundamentally viewed as a PV anomaly than a temperature anomaly. The wave life-cycle begins with an unstable atmosphere which allows growth in PV anomaly amplitude, which in this paradigm, induces anomalies in temperature, geopotential and wind. This view also appeals to the "wave activity" interpretation of eddy features discussed in Section 4f, since PV anomalies relate directly to the EP flux divergence forcing of the zonal mean state. The PV "charge" paradigm, coupled with its generation by instability mechanisms, appears to provide a satisfying physical explanation for the 4-day wave.

5. Summary

This study takes advantage of the vertical coverage and resolution of the Microwave Limb Sounder instrument to analyze 4-day wave features during August - September 1992 and 1993. Strong 4-day wave events are found during both time series. A double-peaked structure is found in temperature with maxima in the upper stratosphere and lower mesosphere and strong vertical phase variation in between, as predicted by the model of Manney and Randel (1993). Four-day eastward period signals are also

evident in MLS ozone, geopotential height and quasi-geostrophic potential vorticity derived from MLS geopotential height. The ozone signal compares well with results from a linear advective-photochemical model which calculates the ozone response to temperature and wind perturbations associated with the 4-day wave. A negative region of quasi-geostrophic potential vorticity gradient coexists with positive EP flux divergence, suggesting instability dynamics plays a role in the wave forcing. The three-dimensional wave structure is shown to resemble the potential vorticity “charge” concept which includes a vertical temperature dipole and anomalies in PV, geopotential and horizontal wind. The PV paradigm, in which temperature, geopotential and wind anomalies are induced by a potential vorticity “charge”, itself the result of instability growth processes, offers an elegant and conceptually satisfying explanation of the physics of the 4-day wave.

Acknowledgments. We want to thank Dr. Jerald Ziemke for assistance in spectral computations and graphics. UKMO data, the result of the diligent work of Dr. Richard Swinbank and Dr. Alan O'Neill, were obtained from the EOS Distributed Active Archive Center (DAAC) at NASA/Goddard Space Flight Center. ISU co-authors are sponsored in part by National Aeronautics and Space Administration Grant NAG 5-2787, Work at the Jet Propulsion Laboratory, California Institute of Technology, was performed under contract with the National Aeronautics and Space Administration,

Figure 1 (a) East-west power spectrum for wave 1 MIS temperature ($T_{retrieved}$) at 72 S for 18 August – 6 September 1992. Dashed contours are at 10, 20 and 30 $K^2 \cdot day$ and solid contours are from 40 to 100 $K^2 \cdot day$ by 10 $K^2 \cdot day$. (b) same as (a) but for temperature derived from MIS geopotential height (T_{geohgt}). (c,d) Same as (a,b) but for 2- 16 September 1993.

Figure 2 Time vs. longitude plot of $T_{retrieved}$ at 64 S and 1 hPa (48 km) filtered for wave 1 and 3-5 day eastward periods for 14 August – 20 September 1992. Dashed contours are from -4 to 0 K by 1 K. Solid contours are from 1 to 4 K by 1 K.

Figure 3 Polar stereographic plots of MIS temperature at 1 hPa filtered for waves 1-3 anti 3--5 day eastward periods for 22-29 August 1992. Latitude circles are at 20, 40, 60 and 80 S with the South Pole at the center. Eastward direction is clockwise with 0 longitude on the positive vertical axis. Dashed contours are from -5 to 0 K by 1 K. Solid contours are from 1 to 8 K by 1 K.

Figure 4 (a) $T_{retrieved}$ amplitude for wave 1 and 3.8 day eastward period for 18 August 6 September 1992. Contours are every 10% interval of the maximum (4.3 K), with values higher than 50% shaded. (b) $T_{retrieved}$ amplitude for wave 1 and 3.75 day eastward period for 2- 16 September 1993. Contours from 9% to 99% of the maximum (5.2 K) by 10%, with values higher than 59% shaded.

Figure 5 (a) $T_{retrieved}$ on a height vs. longitude grid at 72 S filtered for waves 1-3 and 3-5 day eastward rotation period for 22-29 August 1992. (b) Same as (a.) but for 4- 11 September 1993. Contours are every 1 K with solid (dashed) line having positive (negative) values. W (C) refers to warm (cold) anomalies.

Figure 6 (a) MIS geopotential amplitude for wave 1 and 3.8 day eastward period for 18 August 6 September 1992. (b) Same as (a) but for 2- 16 September 1993, 3.75 day period. Contours are at every 10% interval of the maximum with values higher than 50% shaded.

Figure 7 (a) East-west power spectrum for wave 1 MIS ozone (205 GHz) at 72 S for 18 August - 6 September 1992. Contour increment is 0.02 $ppmv^2 \cdot day$. Dashed lines are from 0.02 to 0.08 $ppmv^2 \cdot day$. Solid lines are from 0.10 to 0.20 $ppmv^2 \cdot day$. (b) Same as (a) but for 183 GHz MIS ozone.

Figure 8 (a) Ozone photochemical parameters adapted from Hartmann and Garcia (1979). Γ is the photochemical relaxation frequency and Θ is the linearized ozone response to small temperature perturbations. (b) Wave 1 ozone amplitude (in ppmv) at 60 S for 4-day eastward period from 18 August – 6 September 1992. Lines are model (thick solid), 205 GHz observations (thin solid) and 183 GHz observations (dashed). Error bars were calculated using the procedure described in Section 4d. Large (small) caps show the extent of the 205 (183) GHz ozone error bars.

Figure 9 Unsoothed power spectra for wave 1 at 72 S plotted along with estimated background and 95% confidence level. (a) $T'_{retrieved}$ for the 1993 series at 2 hPa. (b) $T'_{retrieved}$ for the 1993 series at 0.2 hPa. (c) 183 GHz ozone for the 1992 series at 1.0 hPa.

Figure 10 (a) UKMO zonal mean zonal wind (westerly, in Tins-l) averaged from 18 August – 6 September 1992. (b) Quasi-geostrophic potential vorticity gradient, \bar{q}_y , averaged from 18 August - 6 September 1992. Contour increment is $1.0 \times 10^{-11} m^{-1} s^{-1}$ with negative values dashed.

Figure 11 Eliassen-Palm flux diagram calculated for the wave 1, 3.8 day eastward period components of MIS temperature and geopotential height from 18 August - September 1992. Contours are $DP' = \mathbf{V} \cdot \mathbf{F} / (\rho_0 a \cos \phi)$ starting at 0.1 and incremented at 0.4 m/s/day with solid (dashed) line indicating positive (negative) values. Arrows are the Eliassen-Palm flux vectors.

Figure 12 (a) East-west power spectrum for wave 1 quasi-geostrophic potential vorticity derived from MIS geopotential height at 72 S for 18 August - 6 September 1992. (b) Same as (a) but for 2- 16 September 1993. Contours are arbitrary with solid lines representing larger values.

Figure 13 (a) Quasi-geostrophic potential vorticity amplitude for wave 1 and 3.8 day eastward period for 18 August - 6 September 1992. (b) Same as (a) but for 2 - 16 September 1993, 3.75 day period. Contours are at every 10% interval of the maximum with values higher than 50% shaded.

Figure 14 (a) “PV charge” schematic adapted from Bishop and Thorpe (1994) Figs. 1 and 2. The negative PV anomaly induces temperature, geopotential and wind (solid line with arrows) anomalies as shown. Note: the PV, temperature and geopotential anomalies

arc viewed here as a meridional cross-section of the 3-dimensional PV charge, while the circulation is in a horizontal plane which projects out of the page. (b) Amplitude of wave 1, 3.8 day eastward period signal from 18 August -6 September 1992. Hatched region includes PV amplitude greater than 90% of the maximum. Solid lines are perturbation temperature amplitude of 60% to 90%, incremented by 10% of the maximum. Dashed lines are perturbation geopotential amplitude at every 10% of the maximum. Adapted from Figs. 4a, 6a and 13a of this paper.

References

- Andrews, D. G., and M. E. McIntyre, 1978: Generalized Eliassen-Palm and Charney-Drazin theorems for waves on axisymmetric mean flows in compressible atmospheres. *J. Atmos. Sci.*, 35, 175-185.
- Andrews, D. G., and M. E. McIntyre, 1976: Planetary waves in horizontal and vertical shear: the generalized Eliassen-Palm relation and the mean zonal acceleration. *J. Atmos. Sci.*, 33, 2031-2048.
- Barath, P. P., M. C. Chavez, R. E. Cofield, D. A. Flower, M. A. Frerking, M. B. Gram, W. M. Harris, J. R. Holden, R. F. Jarnot, W. G. Kloczeman, G. J. Klose, G. K. Lau, M. S. Loo, B. J. Maddison, R. J. Mattauch, R. P. McKinney, G. E. Peckham, H. M. Pickett, G. Siebes, F. S. Soltis, R. A. Suttie, J. A. Tarsala, J. W. Waters, and W. J. Wilson, 1993: The upper atmosphere research satellite microwave limb sounder instrument, *J. Geophys. Res.*, 98, 10,751-10,762.
- Bishop, C. A., and A. J. Thorpe, 1994: Potential vorticity and the electrostatics analogy: quasi-geostrophic theory. *Q. J. R. Meteorol. Soc.*, 120, 713-731.
- Edmon, H. J. Jr., B. J. Hoskins, and M. E. McIntyre, 1980: Eliassen-Palm cross sections for the troposphere. *J. Atmos. Sci.*, 37, 2600-2616. See also corrigendum, 38, p. 1115.
- Filson, L. S., and L. Froidevaux, 1993: The use of Fourier transforms for asymptotic mapping: applications to the upper atmosphere research satellite microwave limb sounder. *J. Geophys. Res.*, 98, 23039-23049.
- Fishbein, E. F., R. E. Cofield, L. Froidevaux, R. F. Jarnot, T. A. Lungu, W. G. Read, Z. Shippony, J. W. Waters, I. S. McDermid, P. J. McGee, U. Singh, M. Grose, A. Hauchecorne, P. Keckhut, M. E. Gelman, and R. M. Nagatani, 1996: Validation of UARSMLS temperature and pressure measurements. To appear in the *J. Geophys. Res.* special issue on UARS data validation.
- Fraser, G. J., G. Hernandez, and R. W. Smith, 1993: Eastward-moving 2-4 day waves in the winter Antarctic mesosphere. *Geophys. Res. Lett.*, 20, 1547-1550.
- Froidevaux, L., W. G. Read, T. A. Lungu, R. E. Cofield, E. F. Fishbein, D. A. Flower, R. F.

- Jarnot, B.P., Ridenoure, Z., Shippony, J. W., Waters, J. J., Margitan, I. S., McDermid, R. A., Stachnik, G. F., Peckham, G., Braathen, T., Deshler, J., Fishman, D. J., Hofmann, and S. J. Oltmans, 1996: Validation of UARS MLS ozone measurements. To appear in the *J. Geophys. Res.* special issue on UARS data validation.
- Hartmann, D. L., 1983: Barotropic instability of the polar night jet stream. *J. Atmos. Sci.*, 40, 817-835.
- Hartmann, D. L., and R. R. Garcia, 1979: A mechanistic model of ozone transport by planetary waves in the stratosphere. *J. Atmos. Sci.*, 36, 350-364.
- Holton, J. R., 1992: *An introduction to dynamic meteorology; third edition*. Academic Press, Inc., 511 pp.
- Hoskins, B. J., M. F. McIntyre and A. W. Robertson, 1985: On the use and significance of isentropic potential vorticity maps. *Q. J. R. Meteorol. Soc.*, 111, 877-946.
- Lait, L. R., and J. L. Stanford, 1988a: Applications of asymptotic space-time Fourier transform methods to scanning satellite measurements. *J. Atmos. Sci.*, 45, 3784-3799.
- Lait, L. R., and J. L. Stanford, 1988b: Fast, long-lived features in the polar stratosphere. *J. Atmos. Sci.*, 45, 3800-3809.
- Lawrence, B. N., and W. J. Randel, 1996: Variability in the mesosphere observed by the Nimbus 6 PMR. *J. Geophys. Res.*, submitted.
- Lawrence, B. N., G. J. Fraser, R. A. Vincent, and A. Phillips, 1995: The 4-day wave in the Antarctic mesosphere. *J. Geophys. Res.*, 100, 18,899-18,908.
- Madden, R. A., and P. R. Julian, 1971: Detection of a 40-50 day oscillation in the zonal wind in the tropical Pacific. *J. Atmos. Sci.*, 28, 702-708.
- Manney, G. L., R. Swinbank, S. T. Massie, M. F. Gelman, A. J. Miller, R. Nagatani, A. O'Neill, and R. W. Zurek, 1996: Comparison of UKMO and NMC stratospheric analyses during northern and southern winter. To appear in the *J. Geophys. Res.* special issue on UARS data validation.
- Manney, G. L., and W. J. Randel, 1993: Instability at the winter stratopause: a mechanism for the 4-day wave. *J. Atmos. Sci.*, 50, 3928-3938.
- Manney, G. L., 1991: The stratospheric 4-day wave in NMC data. *J. Atmos. Sci.*, 48,

1798-1811.

- Manney, G. L., T. R. Nathan, and J. I. Stanford, 1988: Barotropic stability of realistic stratospheric jets. *J. Atmos. Sci.*, *45*, 2545-2555.
- Matsuno, T., 1970: Vertical propagation of stationary planetary waves in the Northern Hemisphere. *J. Atmos. Sci.*, *27*, 871-883.
- Murakami, M., 1979: Large-scale aspects of deep convective activity over the GATE area. *Mon. Weather Rev.*, *107*, 994-1013.
- Prata, A. J., 1984: The 4-day wave. *J. Atmos. Sci.*, *41*, 150-155.
- Randel, W. J., 1993: Global normal-mode Rossby waves observed in stratospheric ozone data. *J. Atmos. Sci.*, *50*, 406-420.
- Randel, W. J., L. R. Lait, and G. I. Manney, 1992: Dynamics of the 4-day wave: Evidence for instability in the winter upper stratosphere. American Meteorological Society Eighth Conference on Middle Atmosphere, Atlanta, GA January 1992.
- Randel, W. J., and L. R. Lait, 1991: Dynamics of the 4-day wave in the Southern Hemisphere polar stratosphere. *J. Atmos. Sci.*, *48*, 2496-2508.
- Randel, W. J., 1990: Kelvin wave-induced trace constituent oscillations in the equatorial stratosphere. *J. Geophys. Res.*, *95*, 18,641-18,652.
- Ricaud, P., J. de La Noë, B. J. Connor, I. Froidevaux, J. W. Waters, R. S. Harwood, I. A. MacKenzie, and G. E. Peckham, 1996: Diurnal variability of mesospheric ozone as measured by the UARS/MLS instrument: theoretical and ground-based validations. To appear in the *J. Geophys. Res.* special issue on UARS data validation.
- Rood, R. B., and A. R. Douglass, 1985: Interpretation of ozone temperature correlations, I. Theory. *J. Geophys. Res.*, *90*, 5733-5744.
- Salby, M. L., 1982a: Sampling theory for asynoptic satellite observations. Part 1: Space-time spectra, resolution, and aliasing. *J. Atmos. Sci.*, *39*, 2577-2600.
- Salby, M. L., 1982b: Sampling theory for asynoptic satellite observations. Part 1: Fast Fourier synoptic mapping. *J. Atmos. Sci.*, *39*, 2601-2614.
- Thorpe, A. J., and C. H. Bishop, 1995: Potential vorticity and the electrostatics analogy: Fritzel-Rossby formulation. *Q. J. R. Meteorol. Soc.*, *121*, 1477-1495.

- Venne, D. E., and J. L. Stanford, 1982: An observational study of high-latitude stratospheric planetary waves in winter. *J. Atmos. Sci.*, *39*, 1026-1034.
- Venne, D. E., and J. L. Stanford, 1979: Observations of a 4-day temperature wave in the polar winter stratosphere. *J. Atmos. Sci.*, *30*, 2016-2019.
- Ziemke, J. R., and J. L. Stanford, 1990: One-to-two month oscillations in the stratosphere during southern winter. *J. Atmos. Sci.*, *47*, 1778-1793.




Figure 1 (a) East-west power spectrum for wave 1 MLS temperature ($T_{\text{retrieved}}$) at 72 S for 18 August, – 6 September 1992. Dashed contours are at 10, 20 and 30 $K^2 \cdot \text{day}$ and solid contours are from 40 to 100 $K^2 \cdot \text{day}$ by 10 $K^2 \cdot \text{day}$. (b) same as (a) but for temperature derived from MLS geopotential height (T_{geoht}). (c, d) Same as (a, b) but for 2– 16 September 1993,




Figure 2 Time vs. longitude plot of $T_{\text{retrieved}}$ at 64 S and 1 hPa (48 km) filtered for wave 1 and 3-5 day eastward periods for 14 August -- 20 September 1992. Dashed contours are from -4 to 0 K by 1 K. Solid contours are from 1 to 4 K by 1 K.

Tretrieved Waves 1-3 3-5 day (1 hPa)




Figure 3 Polar stereographic plots of MLS temperature at 1 hPa filtered for waves 1-3 and 3-5 day eastward periods for 22-29 August 1992. Latitude circles are at 20, 40, 60 and 80 S with the South Pole at the center. Eastward direction is clockwise with 0 longitude on the positive vertical axis. Dashed contours are from -5 to 0 K by 1 K. Solid contours are from 1 to 8 K by 1 K.

Figure 4 (a) $T_{retrieved}$ amplitude for wave 1 and 3.8 day eastward period for 18 August -6 September 1992. Contours are every 10% interval of the maximum (4.3 K), with values higher than 50% shaded. (b) $T_{retrieved}$ amplitude for wave 1 and 3.75 day eastward period for 2-16 September 1993. Contours from 9% to 99% of the maximum (5.2 K) by 10%, with values higher than 59% shaded.





Figure 5 (a) $T_{\text{retrieved}}$ on a height vs. longitude grid at 72 S filtered for wave-s 1-3 and 3-5 day eastward rotation period for 22-29 August 1992, (b) Same as (a) but for 4 - 11 September 1993. Contours are every 1 K with solid (dashed) line having positive (negative) values. W (C) refers to warm (cold) anomalies.

(a) Geopotential (1992)



(b) Geopotential (1993)





Figure 6 (a) MLS geopotential amplitude for wave 1 and 3.8 day eastward period for 18 August - 6 September 1992. (b) Same as (a) but for 2-16 September 1993, 3.75 day period, Contours are at every 10% interval of the maximum with values higher than 50% shaded.

(a) 205 GHz Ozone



(b) 183 GHz Ozone




Figure 7 (a) East-west power spectrum for wave 1 MLS ozone (205 GHz) at 72 S for 18 August -6 September 1992. Contour increment is $0.02 \text{ ppmv}^2 \cdot \text{day}$. Dashed lines are from 0.02 to $0.08 \text{ ppmv}^2 \cdot \text{day}$. Solid lines are from 0.10 to $0.20 \text{ ppmv}^2 \cdot \text{day}$. (b) Same as (a) but for 183 GHz MLS ozone.




Figure 8 (a) Ozone photochemical parameters adapted from Hartmann and Garcia (1979). Γ is the photochemical relaxation frequency and Θ is the linearized ozone response to small temperature perturbations. (b) Wave 1 ozone amplitude (in ppmv) at 60 S for 4-day eastward period from 18 August – 6 September 1992. Lines are model (thick solid), 205 GHz observations (thin solid) and 183 GHz observations (dashed). Error bars were calculated using the procedure described in Section 4d. Large (small) caps show the extent of the 205 (183) GHz ozone error bars.





Figure 9 Unsmoothed power spectra for wave 1 at 72 S plotted along with estimated background and 95% confidence level. (a) $T_{\text{retrieved}}$ for the 1993 series at 2 hPa. (b) $T_{\text{retrieved}}$ for the 1993 series at 0.2 hPa. (c) 183 GHz ozone for the 1992 series at 1.0 hPa.

(a) UKMO Zonal Wind 18 August -6 September 1992



(b) QGPV Gradient 18 August -6 September 1992




Figure 10 (a) UKMO zonal mean zonal wind (westerly, in $m s^{-1}$) averaged from 18 August -6 September 1992. (b) Quasi-geostrophic potential vorticity gradient, \bar{q}_y , averaged from 18 August - 6 September 1992. Contour increment is $1.0 \times 10^{-11} m^{-1} s^{-1}$ with negative values dashed.




Figure 11 Eliassen-Palm flux diagram calculated for the wave 1, 3.8 day eastward period components of MLS temperature and geopotential height from 18 August – September 1992. Contours are $DF = \nabla \cdot \mathbf{F} / (\rho_o a \cos \phi)$ starting at 0.1 and incremented at 0.4 m/s/day with solid (dashed) line indicating positive (negative) values. Arrows are the Eliassen-Palm flux vectors.





Figure 12 (a) East-west power spectrum for wave 1 quasi-geostrophic potential vorticity derived from MLSgeopotential height at 72 S for 18 August -6 September 1992, (b) Same as (a) but for 2-16 September 1993. Contours are arbitrary with solid lines representing larger values.

(a) Wave 1 PV (1992)



(b) Wave 1 PV (1993)





Figure 13 (a) Quasi-geostrophic potential vorticity amplitude for wave 1 and 3.8 day eastward period for 18 August - 6 September 1992. (b) Same as (a) but for 2 - 16 September 1993, 3.75 day period. Contours are at every 10% interval of the maximum with values higher than 50% shaded.

(a) PV Charge Schematic



(b) 1992 4-Day Wave




Figure 14 (a) "PV charge" schematic adapted from Bishop and Thorpe (1994) Figs. 1 and 2. The negative PV anomaly induces temperature, geopotential and wind (solid line with arrows) anomalies as shown.

Note: the PV, temperature and geopotential anomalies are viewed here as a meridional cross-section of the 3-dimensional circulation is in a horizontal plane which projects out of the page. PV charge, while the

(b) Amplitude of wave 1, 3.8 day eastward period signal from 18 August – 6 September 1992. Hatched region includes PV amplitude greater than 90% of the maximum. Solid lines are perturbation temperature amplitude of 60% to 90%, incremented by 10% of the maximum. Dashed lines are perturbation geopotential amplitude at every 10% of the maximum. Adapted from Figs. 4a, 6a and 13a of this paper.